Corso AINV.
La disautonomia nella pratica
clinica: diagnosi e strategie
terapeutiche

Neuropatia delle piccole fibre:
fisiopatologia, clinica e diagnostica
Vincenzo Provitiera

Neuropatia delle piccole fibre

- Coinvolgimento selettivo o prevalente delle fibre mieliniche di piccolo calibro e amieliniche.

FIBRE SENSITIVE

STIMOLI TERMICI

Caldo

C

Freddo

A δ

Dolore da Caldo

C

Dolore da freddo

A δ

STIMOLI MECCANICI

Tattili

A β

Vibratori

A β

Dolore meccanico

A δ

FIBRE AUTONOMICHE

VISCEROEFFETRICI SECRETOMOTORIE

Noradrenergiche

C

Colinergiche

C

La dimensione del problema

Neurology 2013 Peters MJ et al.

INCIDENZA

11.7 casi/100,000
abitanti/anno

PREVALENZA

52.9 casi/100.000

La dimensione del problema

INCIDENZA

7000 nuovi casi /
anno

PREVALENZA

> 30000

Eziopatogenesi

Metabolic

- ▶ Impaired glucose tolerance
- ▶ Diabetes mellitus
- ▶ Rapid glycaemic control
- ▶ Vitamin B12 deficiency
- ▶ Dyslipidaemia
- ▶ Hypothyroidism
- ▶ Chronic kidney disease

Immune mediated

- ▶ Coeliac disease
- ▶ Sarcoidosis
- ▶ Sjögren's syndrome
- ▶ Rheumatoid arthritis
- ▶ Systemic lupus erythematosus
- ▶ Vasculitis
- ▶ Inflammatory bowel disease
- ▶ Paraneoplastic
- ▶ Monoclonal gammopathy/amyloid

Infections

- ▶ HIV
- ▶ Hepatitis C
- ▶ Influenza

Toxins and drugs

- ▶ Anti-retrovirals
- ▶ Antibiotics—metronidazole, nitrofurantoin, linezolid
- ▶ Chemotherapy—bortezomib
- ▶ Flecainide
- ▶ Statin
- ▶ Alcohol
- ▶ Vitamin B6 toxicity

Hereditary/genetic

- ▶ $Na_v1.7$ mutations
- ▶ $Na_v1.8$ mutations
- ▶ Familial amyloid polyneuropathy
- ▶ Fabry's disease
- ▶ Tangier's disease

Idiopathic

- ▶ Idiopathic small fibre neuropathy
- ▶ Burning mouth syndrome

Caratteristiche cliniche

DISTURBI SENSITIVI

- **Dolore:** peggiora nelle ore notturne
- Intolleranza al contatto con indumenti o lenzuola
- Sindrome delle gambe senza riposo
- Sensazione di intorpidimento, di “fascia” o di “freddo”

- ✓ **Dolore urente**
- ✓ **Freddo doloroso**
- ✓ **Prurito**
- ✓ **Stretta dolorosa**
- ✓ **Dolore lancinante**

le neuropatie delle piccole fibre (SFN)

sintomi e segni_distribuzione

length-dependent
distribution

Non-length dependent small fibre neuropathy/ ganglionopathy

K C Gorson,¹ D N Herrmann,² R Thiagarajan,² T H Brannagan,³ R L Chin,³ L J Kinsella,⁴
A H Ropper⁵

J Neurol Neurosurg Psychiatry 2008;**79**:163–169.

Small fiber neuropathy

Diagnostic criteria

Almeno due criteri tra:

- 1) ipoestesia termica/dolorifica e/o iperalgesia e/o allodinia;
- 2) Anomalie al *quantitative sensory testing* al sito distale
- 3) Ridotta densità di IENF alla gamba

Assenza di segni di coinvolgimento di fibre di grosso calibro (atassia, areflessia, apallestesia...). Normale neurografia.

Small fiber neuropathy Diagnostic criteria

POSSIBILE

Sintomi o segni di coinvolgimento delle piccole fibre o entrambi

PROBABILE

Segni di coinvolgimento delle piccole fibre e normali velocità di conduzione

DEFINITO

Criteri di probabilità

+

Anomalie del QST

o

Riduzione delle fibre epidermiche alla gamba

Why skin biopsy?

- ✓ minimally invasive
- ✓ more sensitive than sural nerve
- ✓ possibility to recognize fibers from their target

Epidermal nerve fibers

✓ Paul Langerhans 1868

“ *First observations of
Epidermal Nerve Fiber* ”

✓ Arthur and Shelley 1959

✓ Wang et al. 1990

“ *Demonstration of Epidermal
Nerve Fiber using PGP* ”

Method standarditation

✓ Kennedy and Crabb 1993

✓ Mc Carthy et al. 1995

Normative data

Bright-field

Immunofluorescence

Lauria G et al. , J Peripher Nerv Syst. 2010

Provitera V et al. , Eur J Neurol. 2016

- ✓ *about 1 fiber/mm more in woman*
- ✓ *0.54 fiber/mm loss for decade*

Bright-field **VS** Immunofluorescence

- ✓ Same diagnostic efficiency
- ✓ FL/BF ratio 2 : 1

Nolano M et al. , J Peripher Nerv Syst. 2015

The role of skin biopsy in differentiating small-fiber neuropathy from ganglionopathy

V. Provitera^a, C. H. Gibbons^a, G. Wendelachate-Crisb^a, V. Donadio^a, D. F. Vitale^a, A. Loaverbruck^a, A. Stancanelli^a, G. Caporaso^a, R. Liguori^a, N. Wang^b, L. Santoro^a, W. R. Kennedy^a and M. Nolano^a

SFN

Thigh

Leg

Table 1 Demographic data

Subject	<i>n</i>	Sex (M/F)	Age (years)
Controls	288	125/163	47.6 ± 13.9
Length-dependent SFN	173	108/65	54.9 ± 14.6
SFSG	141	63/78	51.4 ± 12.9

SFN, small-fiber neuropathy; SFSG, small-fiber sensory ganglionopathy. Data are given as mean ± SD.

ENF Leg/thigh ratio = 0.67 ± 14
 5° percentile cutoff = 0.48
 Sensitivity: 0.80
 Specificity: 0,61

Adjunctive criteria to increase skin biopsy diagnostic efficiency

QUANTITATIVE SENSORY TESTING (QST)

- Soglie Tattili:

MONOFILAMENTI CALIBRATI DI SEMMES WEINSTEIN

- Pinprick:

MONOFILAMENTO CALIBRATO CON PUNTA METALLICA DI 50 MICRON

- Soglie Termiche:

CASE IV - 4-2-1 STEPPING ALGORITHM
TSA 2001 - METODO DEI LIMITI – METODO DEI LIVELLI

DFNS Sensory Profile

- Cold detection threshold (CDT)
- Warm detection threshold (WDT)
- Thermal sensory limen (TSL)
- Paradoxical heat sensations (PHS)
- Cold pain threshold (CPT)
- Heat pain threshold (HPT)
- Tactile detection thresholds (MDT)
- Mechanical pain threshold (MPT)
- Mechanical pain sensitivity (MPS)
- Dynamic mechanical allodynia (DMA)
- Wind-up ratio (WUR)
- Vibration detection threshold (VDT)
- Pressure pain threshold (PPT)

THE RICH SUPPLY OF AUTONOMIC NERVES IN THE DEEPER CUTANEOUS LAYERS

Caratteristiche cliniche

SINTOMI DI DISFUNZIONE AUTONOMICA:

- Ipo o iperidrosi, diarrea o stipsi, gastroparesi
- Sintomi ortostatici
- Incontinenza o ritenzione urinaria
- Visione offuscata, arrossamenti facciali transitori
- Disfunzione sessuale, disturbi vasomotori acrali

Semiquantitative assessment

4

3

2

1

Il controllo del flusso sanguigno nella cute

VASOCOSTRIZIONE

Riduzione
del flusso

Conservazione
termica

VASODILATAZIONE

Aumento
del flusso

Dispersione
termica

VASOCOSTRIZIONE

- Freddo
- Respiro profondo
- Stress emotivo
- Stimoli di arousal
- Cambi di posizioni degli arti

VASODILATAZIONE

- Aumento temp. corporea
- Riscaldamento locale
- Dolore

TRASMISSIONE
ADRENERGICA
alpha1 e alpha2

CO-TRASMISSIONE
COLINERGICA
Ach, NO, neuropeptidi

FLUSSIMETRIA LASER DOPPLER

- Trasmissione di raggio laser
- Registrazione della variazione in lunghezza d'onda rispetto al raggio incidente

Variazione
del flusso

Maggiore
wL shift

IL FLUSSIMETRO LASER DOPPLER

- UNITA' CENTRALE
- SONDE A FIBRA OTTICA
- MODULI PER STIMOLAZIONE TERMICA IONOFRESI
- SOFTWARE PER REGISTRAZIONE E ANALISI DEL SEGNALE

IL SEGNALE LASER DOPPLER

- Campo d'azione 1-1,5 mm di profondità
- Registrazione della variazione di flusso in unità convenzionali di perfusione

PRINCIPALI PROTOCOLLI

FUNZIONE

STIMOLO

RISULTATO

VASODILATAZIONE

STIMOLO TERMICO

INCREMENTO

VASODILATAZIONE

ACh Ionoforesi

INCREMENTO

VASOCOSTRIZIONE

AROUSAL/RESPIRO

RIDUZIONE

VASOCOSTRIZIONE

POSTURA 3-5'

RIDUZIONE

PROTOCOLLO DI STUDIO

QPART (Quantitative pilomotor axon reflex test)

- Fenilefrina 1% per ionoforesi (0.5 mA per 5')
- Applicazione calco in silicone (10'')
- Applicazione di toner per aumentare il contrasto
- Scansione del calco
- Analisi quantitativa

RISULTATI NELLO STUDIO DI VALIDAZIONE

Assenza di risposta per ionofresi con soluzione salina

Assenza di differenze side-to-side in termini di

- 1) densità di impronte nell'area di stimolazione diretta e riflessa
- 2) Estensione dell'area di piloerezione riflessa nella zona circostante l'area stimolata

	Area stimolata dx	Area stimolata sx	Riflesso dx	Riflesso sx
Densità impronte	13.4 $\pm$ 2.7	15.0 $\pm$ 2.0	66.0 $\pm$ 13.9	62.0 $\pm$ 11.3
Area attivazione (cm ²)	-	-	51.0 $\pm$ 3.2	51.5 $\pm$ 3.2

RISULTATI NELLO STUDIO DI VALIDAZIONE

Latenza risposta diretta massimale 10 secondi

Latenza risposta riflessa massimale 600 secondi

RISULTATI NELLO STUDIO DI VALIDAZIONE

Pretrattamento con gel di lidocaina

- 1) Assenza di differenze nella reazione diretta
- 2) Marcata riduzione della risposta riflessa (neuromediata)

QPART è l'unico test validato per la quantizzazione della funzione pilomotoria.

ma...

È stato testato solo in piccole popolazioni di pazienti e mancano dati normative su casistiche estese

Stati emotivi e una temperatura ambientale non controllata possono condizionare il test

La Giunzione sudomotoria

Quantitative sudomotor axon-reflex test - QSART

Ach per
ionofresi

QSART - risultati

Misurazione delle variazioni di umidità nel tempo

P.A. Low / Clinical Neurophysiology 115 (2004) 1506–1513

Parametri

Area sotto la curva

Massima
produzione

Latenza

Silastic imprint test - SIT

Pilocarpina per ionoforesi

Valutazione distribuzione e densità delle ghiandole attivate

Dynmic sweat test

DST - PARAMETRI VALUTATI

- 1. Numero di ghiandole attivate per cm^2
- 2. Sudore prodotto per ghiandola per minuto (nl/min)
- 3. Sudore prodotto per cm^2 per minuto

ORIGINAL COMMUNICATION

Diagnostic value of blood tests for occult causes of initially idiopathic small-fiber polyneuropathy

Magdalena Lang¹ · Roi Treister¹ · Anne Louise Oakland

- 213 pazienti con diagnosi di sfn idiopatica definitiva

Table 2 Prevalence of abnormal test results (ABTR) in the sSMFN cohort and in comparison populations

Test (definition of abnormal result)	Medical condition tested for	Prevalence of ABTR in sample (n)	Population prevalence of ABTR and source of population data
A1C (high)	Hypercortisolism [24]	44.6% (83)	Not evaluated due to positive predictive value = 0
ESR (high)	Inflammation/infection [32, 43]	28.0% (53)	3.0% in Norway [30]
ANA (≥ 1:160)	Lupus/chronic disease [43]	27.3% (53)	8.9% in Brazil [21]
2-hr OGTT value for prediabetes (140–149 mg/dL)	Impaired glucose tolerance (prediabetes) [3]	23.0% (40)	44.9% in US adults 45–64y from A1C, FPG, or 2-hr OGTT value NHANES [40]
Fasting plasma glucose for prediabetes (100–125 mg/dL)	Impaired fasting plasma glucose (prediabetes) [3]	23.0% (40)	40.9% in US adults 45–64y from A1C, FPG, or 2-hr OGTT value NHANES [40]
Triglycerides (high)	Hypertriacylglyceridemia [28]	24.7% (47)	30% NHANES [40]
Complement C4 (low)	Inflammation/vasculitis [43]	15.7% (31)	10.4% WHR [31]
Liver ALT/ALP (high)	Fatty liver, alcoholism, hepatitis [3]	14.8% (30)	10% NHANES [29]
A1C for prediabetes (≥ 5.7% < 6.5)	Recent hyperglycemia (prediabetes)	14.7% (30)	40.9% in US adults 45–64y from A1C, FPG, or 2-hr OGTT value [40]
C-reactive protein (high)	Injury/inflammation [25]	12.6% (25)	7.7% WHR [30]
Complement C3 (low)	Autoimmunity/vasculitis [43]	11.0% (21)	2.3% WHR [31]
AntiRo52-A	Sjögren's syndrome [49, 56]	8.2% (16)	0.7% WHR [31], 3.9% NHANES [54]
AntiRo52-B	Sjögren's syndrome [49, 56]	9.2% (18)	1.2% WHR [31], 2.4% NHANES [54]
Lysine (Lp) Western blot	Lysine disease [25]	8.3% (16)	No data found on normal-kidney patients
A1C for diabetes (≥ 6.5%)	Recent hyperglycemia/diabetes [40]	5.3% (10)	5.9% (adults) by A1C or OGTT age 45–64 NHANES [40]
Thyroid stimulating hormone (TSH) (high)	Hyperthyroidism [3]	4.1% (14)	0.3% NHANES [27]
SPEP/PEP	Monoclonal gammopathy [74]	3.9% (12)	3.2% for age = 50y [23]
IgA TTG antibody (high)	Celiac sprue [9]	3.3% (10)	0.3–0.8% U.S. estimate [20]
Creatinin (high)	Renal disease, false [67]	2.3% (8)	No data found
Thyroid stimulating hormone (TSH) (low)	Hypothyroidism [97]	2.1% (8)	0.3% NHANES [27]
Folate (low)	Folate deficiency [33]	2.0% (8)	0.1% [44]
Vitamin B12 (low)	Vitamin B12 deficiency [68]	1.3% (5)	0.9% [23]
Hepatitis C antibodies	Hepatitis C [10]	1.1% (4)	1.6% NHANES [40]
Fasting glucose for diabetes including OGTT (≥ 126 mg/dL)	Diabetes mellitus [7]	0.0% (0)	5.8% (adults) by A1C or OGTT age 45–64 NHANES [40]
2-hr value from OGTT for diabetes (≥ 200 mg/dL)	Diabetes mellitus [5]	0.0% (0)	5.8% (adults) by A1C or OGTT age 45–64 NHANES [40]

Prevalenza nella popolazione generale vs. prevalenza di test positivi nella popolazione di iSFN

TEST	DISEASE	PERCENT OF POSITIVE TEST IN STUDY POPULATION	PREVALENCE IN GENERAL POPULATION
ANA ($\geq$ 1:160)	Lupus/rheumatic disease	27.5%	8.9%
AntiRo/SS-A	Sjogren's syndrome	9.2%	0.7%
AntiLa/SS-B		9.2%	1.2%
TSH (high)	Hyperthyroidism	4.1%	0.5%
TSH (low)	Hypothyroidism	2.1%	0.3%
IgA TTG antibody (high)	Celiac sprue	3.5%	0.5-1.0%

Prevalenza nella popolazione generale vs. prevalenza di test positivi nella popolazione di iSFN

TEST	DISEASE	PERCENT OF POSITIVE TEST IN STUDY POPULATION	PREVALENCE IN GENERAL POPULATION
C-reactive protein (high)	Injury/inflammation	12.6%	7.1%
Complement C4 (low)	Inflammation/vasculitis	15.7%	10.4%
Liver AST/ALT (high)	Fatty liver, alcoholism, hepatitis	14.8%	10%
SPEP/IFIX	Monoclonal gammopathy	3.9%	3.2%

Prevalenza nella popolazione generale vs. prevalenza di test positivi nella popolazione di iSFN

TEST	DISEASE	PERCENT OF POSITIVE TEST IN STUDY POPULATION	PREVALENCE IN GENERAL POPULATION
HbA1C ($\geq 5.7\%$, <6.5)	prediabetes	14.7%	44.9%
HbA1C ($\geq 6.5\%$)	diabetes	5.5%	5.8%
Vitamin B12 (low)	Vitamin B12 deficiency	1.5%	3.8%
Hepatitis C antibodies	Hepatitis C	1.1%	1.6%
ACE (high)	Sarcoidosis	44.6%	Not evaluated due to positive predictive value = 0

- ANA ($\geq 1:160$)
- AntiRo/SS-A
AntiLa/SS-B
- TSH (high)
- TSH (low)
- IgA TTG
antibody (high)

- C-reactive
protein (high)
- Complement C4
(low)
- Liver AST/ALT
(high)
- SPEP/IFIX

- HbA1C ($\geq 5.7\%$,
<6.5)
- HbA1C ($\geq 6.5\%$)
- Vitamin B12
(low)
- Hepatitis C
antibodies
- ACE (high)

Corso AINV.
La disautonomia nella pratica
clinica: diagnosi e strategie
terapeutiche

Grazie!